

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Sulfolobus DNA Polymerase IV

Catalog Number: M0327S Concentration: 2,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10

nmol of dNTP into acid insoluble material in 30 minutes at 55°C.

Packaging Lot Number: 10225940
Expiration Date: 11/2025
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0327S/L v2.0

Sulfolobus DNA Polymerase IV Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0327SVIAL	Sulfolobus DNA Polymerase IV	10215726	Pass	
B9004SVIAL	ThermoPol® Reaction Buffer Pack	10215580	Pass	

Assay Name/Specification	Lot # 10225940
Endonuclease Activity (Nicking) A 50 μl reaction in ThermoPol® Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 units of Sulfolobus DNA Polymerase IV incubated for 4 hours at 37°C and 55°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 20 units of Sulfolobus DNA Polymerase IV incubated for 4 hours at 37°C and 55°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 2 units of Sulfolobus DNA Polymerase IV incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE)	Pass



M0327S / Lot: 10225940

Page 1 of 2

Assay Name/Specification	Lot # 10225940
Sulfolobus DNA Polymerase IV is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
Single Stranded DNase Activity (FAM-Labeled Oligo) A 20 µl reaction in ThermoPol® Reaction Buffer containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 20 units of Sulfolobus DNA Polymerase IV incubated for 30 minutes at 37°C and 55°C yields <10% degradation as determined by capillary electrophoresis.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 2 units of Sulfolobus DNA Polymerase IV is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

30 Nov 2023

Michael Tonello

Packaging Quality Control Inspector

30 Jan 2024



M0327S / Lot: 10225940 Page 2 of 2