

## New England Biolabs Certificate of Analysis


**Product Name:** *Bst DNA Polymerase, Full Length*  
**Catalog Number:** M0328S  
**Concentration:** 5,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 65°C.  
**Lot Number:** 10055642  
**Expiration Date:** 07/2021  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 % Triton®X-100, 50 % Glycerol, (pH 7.1 @ 25°C)  
**Specification Version:** PS-M0328S/L v1.0

Bst DNA Polymerase, Full Length Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0328SVIAL	Bst DNA Polymerase, Full Length	10049393	Pass
B9004SVIAL	ThermoPol® Reaction Buffer Pack	10041932	Pass

Assay Name/Specification	Lot # 10055642
<p><b>RNase Activity (Extended Digestion)</b>            A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Bst DNA Polymerase, Full Length is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p><b>qPCR DNA Contamination (E. coli Genomic)</b>            A minimum of 5 units of Bst DNA Polymerase, Full Length is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p><b>Protein Purity Assay (SDS-PAGE)</b>            Bst DNA Polymerase, Full Length is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p><b>Phosphatase Activity (pNPP)</b>            A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl<sub>2</sub> containing 2.5 mM</p>	Pass

Assay Name/Specification	Lot # 10055642
<p>p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Bst DNA Polymerase, Full Length incubated for 4 hours at 37°C yields &lt;0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 units of Bst DNA Polymerase, Full Length incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Bst DNA Polymerase, Full Length incubated for 4 hours at 37°C and 65°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.




---

Doreen Duquette  
Production Scientist  
18 Apr 2019




---

Josh Hersey  
Packaging Quality Control Inspector  
12 Sep 2019