

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Exonuclease V (RecBCD)

Catalog Number: M0345S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to produce 1

nmol of acid-soluble deoxyribonucleotide from double-stranded DNA in

30 minutes at 37°C in a total reaction volume of 50 μl.

Lot Number: 10051887
Expiration Date: 08/2021
Storage Temperature: -20°C

Storage Conditions: 50 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

0.1% Triton®X-100, (pH 7.5 @ 25°C)

Specification Version: PS-M0345S/L v1.0

Exonuclease V (RecBCD) Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
P0756SVIAL	Adenosine 5'-Triphosphate (ATP)	10045567	Pass	
M0345SVIAL	Exonuclease V (RecBCD)	10051890	Pass	
B7004SVIAL	NEBuffer™ 4	10043904	Pass	

Assay Name/Specification	Lot # 10051887
Endonuclease Activity (Nicked Double-Stranded DNA) A 50 μl reaction in NEBuffer 4 supplemented with 1 mM ATP containing 1 μg of nicked PhiX174 RF II DNA and a minimum of 50 units of Exonuclease V (RecBCD) incubated for 4 hours at 37°C results in <10% loss in PhiX174 RF II DNA as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 4 supplemented with 1 mM ATP containing 1 μg of supercoiled PhiX174 RF I DNA and a minimum of 100 units of Exonuclease V (RecBCD) incubated for 4 hours at 37°C results in <10% loss in supercoiled DNA as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Exonuclease V (RecBCD) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion)	Pass



M0345S / Lot: 10051887

Page 1 of 2

Assay Name/Specification	Lot # 10051887
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	
and a minimum of 10 units of Exonuclease V (RecBCD) is incubated at 37°C. After	
incubation for 4 hours, >90% of the substrate RNA remains intact as determined by	
gel electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

Alicia Bielik Production Scientist

30 May 2019

Michael Tonello

Packaging Quality Control Inspector

21 Aug 2019

