

New England Biolabs Certificate of Analysis

Product Name: T4 RNA Ligase 2, truncated K227Q
Catalog Number: M0351L
Concentration: 200,000 U/ml
Unit Definition: 200 units is defined as the amount of enzyme required to give 80% ligation of a 31-mer RNA to the pre-adenylated end of a 17-mer DNA in a total reaction volume of 20 µl in 1 hour at 25°C.
Lot Number: 10008154
Expiration Date: 04/2020
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 100 mM NaCl, 0.1 mM DTT, 0.1 mM EDTA, 50% Glycerol, (pH 7.5 @ 25°C)
Specification Version: PS-M0351S/L v1.0

T4 RNA Ligase 2, truncated K227Q Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0351LVIAL	T4 RNA Ligase 2, truncated K227Q	0011804	Pass
B1004SVIAL	PEG 8000	0151708	Pass
B0216SVIAL	T4 RNA Ligase Reaction Buffer	10008137	Pass

Assay Name/Specification	Lot # 10008154
Protein Purity Assay (SDS-PAGE) T4 RNA Ligase 2, truncated K227Q is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 200 units of T4 RNA Ligase 2, truncated K227Q incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 200 units of T4 RNA Ligase 2, truncated K227Q is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Endonuclease Activity (Nicking)	Pass

Assay Name/Specification	Lot # 10008154
<p>A 50 µl reaction in T4 RNA Ligase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 200 units of T4 RNA Ligase 2, truncated K227Q incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p> <p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in T4 RNA Ligase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of T4 RNA Ligase 2, truncated K227Q incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.



Bo Wu
Production Scientist
17 May 2018



Michael Tonello
Packaging Quality Control Inspector
12 Jun 2018