

New England Biolabs Certificate of Analysis

Product Name: T4 RNA Ligase 2, truncated K227Q
Catalog Number: M0351L
Concentration: 200,000 U/ml
Unit Definition: 200 units is defined as the amount of enzyme required to give 80% ligation of a 31-mer RNA to the pre-adenylated end of a 17-mer DNA in a total reaction volume of 20 µl in 1 hour at 25°C.
Packaging Lot Number: 10249058
Expiration Date: 04/2026
Storage Temperature: -20°C
Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0351S/L v2.0

T4 RNA Ligase 2, truncated K227Q Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0351LVIAL	T4 RNA Ligase 2, truncated K227Q	10238093	Pass
B1004SVIAL	PEG 8000	10218040	Pass
B0216SVIAL	T4 RNA Ligase Reaction Buffer	10239990	Pass

Assay Name/Specification	Lot # 10249058
<p>Endonuclease Activity (Nicking) A 50 µl reaction in T4 RNA Ligase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 200 units of T4 RNA Ligase 2, truncated K227Q incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in T4 RNA Ligase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of T4 RNA Ligase 2, truncated K227Q incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Phosphatase Activity (pNPP) A 200 ul reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 200 units of T4 RNA Ligase 2, truncated K227Q incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass

Assay Name/Specification	Lot # 10249058
<p>Protein Purity Assay (SDS-PAGE) T4 RNA Ligase 2, truncated K227Q is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 200 units of T4 RNA Ligase 2, truncated K227Q is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Bo Wu
Production Scientist
01 May 2024



Josh Hersey
Packaging Quality Control Inspector
03 Jul 2024