

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: mRNA Cap 2'-O-Methyltransferase

Catalog Number: M0366S
Concentration: 50,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to methylate 10

pmoles of 80 nt long capped RNA transcript in 1 hour at 37°C.

Packaging Lot Number: 10132676
Expiration Date: 08/2023
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl , 20 mM Tris-HCl (pH 8.0), 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol , 0.1 % Triton®X-100

Specification Version: PS-M0366S v1.0

mRNA Cap 2'-O-Methyltransferase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0366SVIAL	mRNA Cap 2'-O-Methyltransferase	10117905	Pass	
B9003SVIAL	S-adenosylmethionine (SAM)	10124322	Pass	
B2080AVIAL	10X Capping Buffer	10120094	Pass	

Assay Name/Specification	Lot # 10132676
Protein Purity Assay (SDS-PAGE) mRNA Cap 2'-O-Methyltransferase is ≥ 99% pure as determined by SDS-PAGE analysis	Pass
using Coomassie Blue detection.	
RNase Activity (Extended Digestion)	Pass
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 50 units of mRNA Cap 2'-O-Methyltransferase is incubated at 37°C.	
After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	
Francisco Asticito (Padiaceticito Balacca)	Bass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in Capping Buffer containing 1 μg of a mixture of single and	Pass
double-stranded [ ³H] E. coli DNA and a minimum of 50 units of mRNA Cap 2'-O-Methyltransferase incubated for 4 hours at 37°C releases <0.1% of the total	
radioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in Capping Buffer containing 1 µg of supercoiled PhiX174 DNA and a	



M0366S / Lot: 10132676

Page 1 of 2

Assay Name/Specification	Lot # 10132676
minimum of 50 units of mRNA Cap 2'-O-Methyltransferase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist

28 Dec 2021

Josh Hersey

Packaging Quality Control Inspector

28 Dec 2021



M0366S / Lot: 10132676 Page 2 of 2