

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Bst 3.0 DNA Polymerase
Catalog #:	M0374M
Concentration:	120,000 units/ml
Unit Definition:	One unit is defined at the amount of enzyme that will incorporate 25 nmol of dNTPs into acid insoluble material in 30 minutes at 65° C.
<i>Lot</i> #:	0071712
Assay Date:	12/2017
Expiration Date:	12/2019
Storage Temp:	-20°C
Storage Conditions:	10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.1 % Triton®X-100 , 50 % Glycerol, (pH 7.4 @ 25° C)
Specification Version:	PS-M0374M v1.0
Effective Date:	02 Sep 2015

Assay Name/Specification (minimum release criteria)	Lot #0071712
Endonuclease Activity (Nicking) - A 50 μ l reaction in ThermoPol® Reaction Buffer containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 500 units of <i>Bst</i> 3.0 DNA Polymerase incubated for 4 hours at 65°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) - A 50 μ l reaction in ThermoPol® Reaction Buffer containing 1 μ g of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 500 units of <i>Bst</i> 3.0 DNA Polymerase incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 120 units of <i>Bst</i> 3.0 DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Phosphatase Activity (pNPP) - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 mM <i>p</i> -Nitrophenol Phosphate (pNPP) and a minimum of 100 units <i>Bst</i> 3.0 DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE) - <i>Bst</i> 3.0 DNA Polymerase is \geq 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



M0374M Lot: 0071712 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0071712
qPCR DNA Contamination (<i>E. coli</i> Genomic) - A minimum of 120 units of <i>Bst</i> 3.0 DNA Polymerase is screened for the presence of <i>E. coli</i> genomic DNA using SYBR® Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.	Pass
RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single- stranded RNA and a minimum of 1 μ l of <i>Bst</i> 3.0 DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

Authorized by Melanie Fortier 02 Sep 2015



1llho In

Inspected by Tony Spear-Alfonso 13 Dec 2017