

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: SplintR® Ligase

Catalog Number: M0375L Concentration: 25,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme needed to ligate (to 50%

completion) 2 picomoles of a tripartite FAM-labeled DNA:RNA hybrid substrate in a 20 µl reaction at 25°C in 15 minutes in 1X SplintR®

Ligase Reaction Buffer.

Lot Number: 10040000
Expiration Date: 03/2021
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0375S/L v1.0

| SplintR® Ligase Component List |                                     |            |                      |  |
|--------------------------------|-------------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>         | Component Description               | Lot Number | Individual QC Result |  |
| M0375LVIAL                     | SplintR® Ligase                     | 10040001   | Pass                 |  |
| B0375SVIAL                     | 10X SplintR® Ligase Reaction Buffer | 10020201   | Pass                 |  |

| Assay Name/Specification   | Lot # 10040000 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 μl reaction in SplintR® Ligase Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 125 units of SplintR® Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in SplintR® Ligase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 125 units of SplintR® Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.   | Pass           |
| Protein Purity Assay (SDS-PAGE) SplintR® Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.   | Pass           |
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA  | Pass           |



M0375L / Lot: 10040000

Page 1 of 2

electrophoresis using fluorescent detection.

This product has been tested and shown to be in compliance with all specifications.

Bo Wu

Production Scientist 12 Sep 2018 Michael Tonello

Packaging Quality Control Inspector

02 Apr 2019

