

## New England Biolabs Certificate of Analysis

**Product Name:** Exonuclease VII  
**Catalog Number:** M0379S  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme that will catalyze the release of 1 nmol of acid-soluble nucleotide in a total reaction volume of 50 µl in 30 minutes at 37°C.  
**Packaging Lot Number:** 10080133  
**Expiration Date:** 08/2022  
**Storage Temperature:** -20°C  
**Storage Conditions:** 100 mM NaCl, 50 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.1 % Triton®X-100, (pH 7.5 @ 25°C)  
**Specification Version:** PS-M0379S/L v1.0


Exonuclease VII Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0379SVIAL	Exonuclease VII	10080132	Pass
B0379SVIAL	Exonuclease VII Reaction Buffer	10058706	Pass

Assay Name/Specification	Lot # 10080133
<p><b>Non-Specific DNase Activity (16 Hour)</b>            A 50 µl reaction in NEBuffer 4 containing 1 µg of HaeIII digested PhiX174 RF I DNA and a minimum of 10 units of Exonuclease VII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p><b>Exonuclease Activity (Radioactivity Release, Double Stranded)</b>            A 50 µl reaction in NEBuffer 4 containing 1 µg double stranded [<sup>3</sup>H] E. coli DNA and a minimum of 10 units of Exonuclease VII incubated for 4 hours at 37°C releases &lt;0.5% of the total radioactivity.</p>	Pass
<p><b>Endonuclease Activity (Nicking)</b>            A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of Exonuclease VII incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p><b>Endonuclease Activity (Circular Single Stranded DNA)</b>            A 50 µl reaction in NEBuffer 4 containing 1 µg of M13 single-stranded DNA and a</p>	Pass

Assay Name/Specification	Lot # 10080133
<p>minimum of 10 units of Exonuclease VII incubated for 1 hour at 37°C results in &lt;20% conversion to linear DNA as determined by agarose gel electrophoresis.</p>	
<p><b>Protein Purity Assay (SDS-PAGE)</b> Exonuclease VII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>RNase Activity Assay (4 Hour Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 10 units of Exonuclease VII is incubated at 37°C. After incubation for 4 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 10 units of Exonuclease VII is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



John Greci  
Production Scientist  
16 Sep 2020



Michael Tonello  
Packaging Quality Control Inspector  
16 Sep 2020