

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: WarmStart® RTx Reverse Transcriptase

Catalog Number: M0380S Concentration: 15,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 1

nmol of dTTP into acid-insoluble material in 20 minutes at 50°C.

Lot Number: 10043534
Expiration Date: 05/2021
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0380S/L v2.0

| WarmStart® RTx Reverse Transcriptase Component List |                                      |            |                      |  |
|---|--------------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>                              | Component Description                | Lot Number | Individual QC Result |  |
| M0380SVIAL  | WarmStart® RTx Reverse Transcriptase | 10043507   | Pass                 |  |
| B1003SVIAL  | Magnesium Sulfate (MgSO₄) Solution   | 0021701    | Pass                 |  |
| B0537SVIAL  | Isothermal Amplification Buffer      | 10035085   | Pass                 |  |

| Assay Name/Specification   | Lot # 10043534 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 μl reaction in Isothermal Amplification Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 15 units of RTx Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.  | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in Isothermal Amplification Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 15 units of RTx Reverse Transcriptase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.  | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 2 containing 1 μg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 15 units of WarmStart® RTx Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |



M0380S / Lot: 10043534

Page 1 of 2

| Assay Name/Specification   | Lot # 10043534 |
|--|----------------|
| Protein Purity Assay (SDS-PAGE) RTx Reverse Transcriptase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.   | Pass           |
| <b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 15 units of WarmStart® RTx Reverse Transcriptase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome. | Pass           |
| RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of WarmStart® RTx Reverse Transcriptase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined   | Pass           |

This product has been tested and shown to be in compliance with all specifications.

by gel electrophoresis using fluorescent detection.

Doreen Duquette **Production Scientist** 

21 Feb 2019

Michael Tonello Packaging Quality Control Inspector 07 May 2019



M0380S / Lot: 10043534