

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Cas9 Nuclease, S. pyogenes

Catalog Number:M0386TConcentration: $20 \mu M$ Lot Number:10028147Expiration Date:01/2021Storage Temperature: $-20^{\circ}C$

Storage Conditions: 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0386T/M v1.0

Cas9 Nuclease, S. pyogenes Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0386TVIAL	Cas9 Nuclease, S. pyogenes	10028148	Pass	
B7203SVIAL	NEBuffer™ 3.1	10021113	Pass	

Assay Name/Specification	Lot # 10028147
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 1 pmol of Cas9 Nuclease, S. pyogenes incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 1 pmol of Cas9 Nuclease, S. pyogenes incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (Targeted Digestion) A 20 µl reaction in NEBuffer 3.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM sgRNA, and 100 nM Cas9 Nuclease, S. pyogenes incubated for 1 hour at 37°C results in ≥90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of Cas9 Nuclease, S. pyogenes incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel	Pass



M0386T / Lot: 10028147

Page 1 of 2

Assay Name/Specification	Lot # 10028147
electrophoresis.	
Protein Purity Assay (SDS-PAGE) Cas9 Nuclease, S. pyogenes is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 pmol of Cas9 Nuclease, S. pyogenes is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Tony Spear-Alfonso

Production Scientist 26 Nov 2018

Michael Tonello

Packaging Quality Control Inspector

18 Jan 2019



M0386T / Lot: 10028147

Page 2 of 2