

New England Biolabs Certificate of Analysis

Product Name: *Cas9 Nuclease, S. pyogenes*
Catalog Number: M0386T
Concentration: 20 μ M
Packaging Lot Number: 10243952
Expiration Date: 05/2026
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0386T/M v1.0

Cas9 Nuclease, <i>S. pyogenes</i> Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0386TVIAL	Cas9 Nuclease, <i>S. pyogenes</i>	10241677	Pass
B6003SVIAL	NEBuffer™ r3.1	10227734	Pass

Assay Name/Specification	Lot # 10243952
Endonuclease Activity (Nicking) A 50 μ l reaction in NEBuffer 3.1 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 1 pmol of Cas9 Nuclease, <i>S. pyogenes</i> incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μ l reaction in NEBuffer 3.1 containing 1 μ g of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 1 pmol of Cas9 Nuclease, <i>S. pyogenes</i> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (Targeted Digestion) A 20 μ l reaction in NEBuffer 3.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM sgRNA, and 100 nM Cas9 Nuclease, <i>S. pyogenes</i> incubated for 1 hour at 37°C results in \geq 90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μ l reaction in NEBuffer 3.1 containing 1 μ g of Lambda DNA and a minimum of 1 pmol of Cas9 Nuclease, <i>S. pyogenes</i> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel	Pass

Assay Name/Specification	Lot # 10243952
electrophoresis.	
Protein Purity Assay (SDS-PAGE) Cas9 Nuclease, <i>S. pyogenes</i> is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 pmol of Cas9 Nuclease, <i>S. pyogenes</i> is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



 Jessica Cane
 Production Scientist
 15 May 2024



 Talia Monkiewicz
 Packaging Quality Control Inspector
 30 May 2024