

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Apyrase
Catalog Number:	M0398S
Concentration:	500 U/ml
Unit Definition:	One unit is defined as the amount of enzyme that catalyses the release of 1 μmol of inorganic phosphate from ATP in 1 minute at 30°C in a total reaction of 50 μl.
Packaging Lot Number:	10162925
Expiration Date:	03/2024
Storage Temperature:	-20°C
Storage Conditions:	20 mM MES, 50 mM NaCl, 1 mM DTT, 0.1 mM CaCl2, 0.1 % Tween® 20, 50 % Glycerol, (pH 6.5 @ 25°C)
Specification Version:	PS-M0398S/L v1.0

Apyrase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0398SVIAL	Apyrase	10162924	Pass	
B0398SVIAL	Apyrase Reaction Buffer	10149806	Pass	

Assay Name/Specification	Lot # 10162925
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 p-Nitrophenyl Phosphate (pNPP) and a minimum of 5 units of Apyrase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE) Apyrase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Apyrase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μI reaction in NEBuffer 4 containing 1 μg of PhiX174-HaeIII DNA and a minimum	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10162925
of 5 units of Apyrase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in Apyrase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 5 units of Apyrase incubated for 4 hours at 30°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in Apyrase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 units of Apyrase incubated for 4 hours at 30°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

barrel Speulding

Darcie Spaulding Production Scientist 09 Sep 2022

Michae

Michael Tonello Packaging Quality Control Inspector 08 Nov 2022

