

New England Biolabs Certificate of Analysis

Product Name: Bst 2.0 WarmStart® DNA Polymerase (Glycerol-free)
Catalog Number: M0402L
Concentration: 120,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that will incorporate 25 nmol of dNTP into acid insoluble material in 30 minutes at 65°C.
Packaging Lot Number: 10259658
Expiration Date: 10/2026
Storage Temperature: -80°C
Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.1 % Triton® X-100, (pH 7.1 @ 25°C)
Specification Version: PS-M0402L v1.0

Bst 2.0 WarmStart® DNA Polymerase (Glycerol-free) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0402LVIAL	Bst 2.0 WarmStart® DNA Polymerase (Glycerol-free)	10259659	Pass
B1714SVIAL	Isothermal Amplification Buffer (Lyo-compatible)	10247890	Pass

Assay Name/Specification	Lot # 10259658
<p>Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 500 units of Bst 2.0® DNA Polymerase (Glycerol-Free) incubated for 4 hours at 65°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 500 units of Bst 2.0® DNA Polymerase (Glycerol-Free) incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Functional Testing (DNA-LAMP) A 25 µl LAMP reaction with 8 units of Bst 2.0 WarmStart® DNA Polymerase (Glycerol-free), 10 ng of genomic DNA and 1X LAMP fluorescent dye results in a threshold time of ≤ 20 minutes as determined by fluorescent detection.</p>	Pass

Assay Name/Specification	Lot # 10259658
<p>Functional Testing (RT-LAMP) A 25 µl RT-LAMP reaction with 8 units of Bst 2.0 WarmStart® DNA Polymerase (Glycerol-free), 10 ng of genomic RNA and 1X LAMP fluorescent dye results in a threshold time of ≤ 20 minutes as determined by fluorescent detection.</p>	Pass
<p>Inhibition of Primer Extension (Hot Start) A 50 µl reaction in Isothermal Amplification Buffer containing 6 mM MgSO₄ and 1.4 mM dNTPs in the presence of 1.6 µM of a fluorescent internally labeled oligonucleotide and a minimum of 16 units of Bst 2.0 WarmStart® DNA Polymerase (Glycerol-Free) incubated for 2 hours at 25°C yields <5% extension as determined by capillary electrophoresis.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 120 units of Bst 2.0 WarmStart® DNA Polymerase (Glycerol-free) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Bst 2.0® DNA Polymerase (Glycerol-Free) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) Bst 2.0® DNA Polymerase (Glycerol-Free) is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 120 units of Bst 2.0 WarmStart® DNA Polymerase (Glycerol-Free) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 120 units of Bst 2.0 WarmStart® DNA Polymerase (Glycerol-Free) is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Trinh Nguyen
Production Scientist
29 Oct 2024



Josh Hersey
Packaging Quality Control Inspector
30 Oct 2024