

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer  |
|------------------------|---|
| Catalog Number:        | M0488L  |
| Concentration:         | 2 X Concentrate   |
| Packaging Lot Number:  | 10146688  |
| Expiration Date:       | 12/2023   |
| Storage Temperature:   | -20°C   |
| Specification Version: | PS-M0488S/L v2.0  |
| Composition (1X):      | 20 mM Tris-HCl (pH 8.9 @ 25°C), 22 mM KCl, 22 mM NH4Cl, 1.8 mM<br>MgCl2, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 %<br>Glycerol, 0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 1 X Xylene<br>cyanol, 1 X Tartrazine, 25 units/ml OneTaq® Hot Start DNA Polymerase |

| OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer Component List |  |            |                      |  |
|---|--|------------|----------------------|--|
| NEB Part Number   | Component Description  | Lot Number | Individual QC Result |  |
| M0488SVIAL  | OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer | 10132090   | Pass                 |  |

| Assay Name/Specification   | Lot # 10146688 |
|--|----------------|
| <b>Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation)</b><br>A 50 µl primer extension assay in ThermoPol® Reaction Buffer in the presence of 200<br>µM dNTPs including [ <sup>3</sup> H]-dTTP, containing 15 nM primed single-stranded M13mp18 with<br>2.5 units of OneTaq® Hot Start DNA Polymerase incubated for 16 hours at 25°C yields<br>>95% inhibition when compared to a non-hot start control reaction.  | Pass           |
| PCR Amplification (Hot Start 2 kb Lambda DNA)<br>A 25 $\mu$ I reaction in OneTaq® Standard Reaction Buffer in the presence of 200 $\mu$ M dNTPs<br>and 0.2 $\mu$ M primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with<br>0.625 units of OneTaq® Hot Start DNA Polymerase for 30 cycles of PCR amplification<br>results in an increase in yield of the 2 kb Lambda product and a decrease in<br>non-specific genomic bands when compared to a non-hot start control reaction. | Pass           |
| <b>PCR Amplification (5 kb Lambda, Master Mix)</b><br>A 25 μl reaction in 1X OneTaq® Hot Start Quick-Load® Master Mix with Standard Buffer<br>and 0.2 μM primers containing 5 ng Lambda DNA for 25 cycles of PCR amplification<br>results in the expected 5 kb product.  | Pass           |
| <b>RNase Activity (Extended Digestion)</b><br>A 10 μI reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA  | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification  | Lot # 10146688 |
|---|----------------|
| and a minimum of 1 $\mu$ I of OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.   |                |
| <b>Non-Specific DNase Activity (16 hour, Buffer)</b><br>A 50 µl reaction in 1X OneTaq® Hot Start Quick-Load® Master Mix with Standard Buffer<br>containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII<br>DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable<br>nuclease degradation as determined by agarose gel electrophoresis. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

vistie Vayanez

Christie Vazquez Production Scientist 29 Mar 2022

Michae

Michael Tonello Packaging Quality Control Inspector 29 Mar 2022

