

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: OneTag® Hot Start Quick-Load® 2X Master Mix with Standard Buffer

Catalog Number: M0488L

Concentration: 2 X Concentrate

Packaging Lot Number: 10207808
Expiration Date: 02/2025
Storage Temperature: -20°C

Specification Version: PS-M0488S/L v2.0

Composition (1X): 20 mM Tris-HCl (pH 8.9 @ 25°C), 22 mM KCl, 22 mM NH4Cl, 1.8 mM

MgCl2, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 1 X Xylene cyanol, 1 X Tartrazine, 25 units/ml OneTaq® Hot Start DNA Polymerase

| OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer Component List | | | |
|---|--|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| M0488SVIAL | OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer | 10189197 | Pass |

| Assay Name/Specification | Lot # 10207808 |
|--|----------------|
| Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation) A 50 µl primer extension assay in ThermoPol® Reaction Buffer in the presence of 200 µM dNTPs including [³H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 2.5 units of OneTaq® Hot Start DNA Polymerase incubated for 16 hours at 25°C yields >95% inhibition when compared to a non-hot start control reaction. | Pass |
| Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X OneTaq® Hot Start Quick-Load® Master Mix with Standard Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| PCR Amplification (5 kb Lambda, Master Mix) A 25 µl reaction in 1X OneTaq® Hot Start Quick-Load® Master Mix with Standard Buffer and 0.2 µM primers containing 5 ng Lambda DNA for 25 cycles of PCR amplification results in the expected 5 kb product. | Pass |
| PCR Amplification (Hot Start 2 kb Lambda DNA) A 25 μl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with | Pass |



M0488L / Lot: 10207808

Page 1 of 2

| Assay Name/Specification | Lot # 10207808 |
|---|----------------|
| 0.625 units of OneTaq® Hot Start DNA Polymerase for 30 cycles of PCR amplification results in an increase in yield of the 2 kb Lambda product and a decrease in non-specific genomic bands when compared to a non-hot start control reaction. | |
| RNase Activity (Extended Digestion) | Pass |
| A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA | |
| and a minimum of 1 µl of OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard | |
| Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA | |

This product has been tested and shown to be in compliance with all specifications.

remains intact as determined by gel electrophoresis using fluorescent detection.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lea Antonpoulos | Reduction Scientist

07 Jun 2023

Michael Tonello

Packaging Quality Control Inspector

25 Sep 2023

