

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: OneTag® Hot Start Quick-Load® 2X Master Mix with GC Buffer

Catalog Number: M0489L

Concentration: 2 X Concentrate

Lot Number: 10053241
Expiration Date: 08/2021
Storage Temperature: -20°C

Specification Version: PS-M0489S/L v1.0

Composition (1X): 80 mM Tris-SO4 (pH 9.2 @ 25°C), 20 mM (NH4)2SO4, 2 mM MgSO4, 0.2 mM

dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 5 % DMSO,

0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 1 X Xylene cyanol, 1 X

Tartrazine, 25 units/ml OneTaq® Hot Start DNA Polymerase

| OneTaq® Hot Start Quick-Load® 2X Master Mix with GC Buffer Component List |  |            |                      |  |
|---|--|------------|----------------------|--|
| <b>NEB Part Number</b>  | Component Description                                      | Lot Number | Individual QC Result |  |
| M0489SVIAL  | OneTaq® Hot Start Quick-Load® 2X Master Mix with GC Buffer | 10050006   | Pass                 |  |
| B9026AVIAL  | OneTaq® High GC Enhancer                                   | 10031487   | Pass                 |  |

| Assay Name/Specification   | Lot # 10053241 |
|--|----------------|
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of OneTaq® Hot Start Quick-Load® 2X Master Mix with GC Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.   | Pass           |
| PCR Amplification (Hot Start 2 kb Lambda DNA) A 25 μl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with 0.625 units of OneTaq® Hot Start DNA Polymerase for 30 cycles of PCR amplification results in an increase in yield of the 2 kb Lambda product and a decrease in non-specific genomic bands when compared to a non-hot start control reaction. | Pass           |
| PCR Amplification (Enhancer Dependent, >70% GC-rich, Master Mix) A 25 μl reaction in 1X OneTaq® Hot Start Quick-Load® Master Mix with GC Buffer and 20% OneTaq® High GC Enhancer in the presence of 0.2 μM primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the enhancer-dependent production of the 627 bp product.   | Pass           |



M0489L / Lot: 10053241

Page 1 of 2

| Assay Name/Specification  | Lot # 10053241 |
|---|----------------|
| PCR Amplification (Buffer Dependent, >65% GC-rich, Master Mix) A 25 μl reaction in 1X OneTaq® Hot Start Quick-Load® Master Mix with GC Buffer and 0.2 μM primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the buffer-dependent production of the 737 bp product.  | Pass           |
| Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X OneTaq® Hot Start Quick-Load® Master Mix with GC Buffer containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.  | Pass           |
| Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation)  A 50 µl primer extension assay in ThermoPol® Reaction Buffer in the presence of 200 µM dNTPs including [ ³H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 2.5 units of OneTaq® Hot Start DNA Polymerase incubated for 16 hours at 25°C yields >95% inhibition when compared to a non-hot start control reaction. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

Production Scientist 13 Feb 2019 Michael Tonello

Packaging Quality Control Inspector

19 Aug 2019

