

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Q5® Hot Start High-Fidelity DNA Polymerase
Catalog Number:	M0493S
Concentration:	2,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 74°C.
Packaging Lot Number:	10110303
Expiration Date:	02/2023
Storage Temperature:	-20°C
Storage Conditions:	Proprietary
Specification Version:	PS-M0493S/L v1.0

Q5® Hot Start High-Fidelity DNA Polymerase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0493SVIAL	Q5® Hot Start High-Fidelity DNA Polymerase	10102369	Pass	
B9028AVIAL	Q5® High GC Enhancer	10107124	Pass	
B9027SVIAL	Q5® Reaction Buffer Pack	10107123	Pass	

Assay Name/Specification	Lot # 10110303
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 2 units of Q5® High-Fidelity DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is $\leq$ 1 E. coli genome.	Pass
Protein Purity Assay (SDS-PAGE) Q5® High-Fidelity DNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Q5® High-Fidelity DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
PCR Amplification (Hot Start, Human Genomic DNA) A 50 µI reaction in Q5® Reaction Buffer plus Q5® High GC Enhancer in the presence of	Pass





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10110303
200 $\mu$ M dNTPs and 0.5 $\mu$ M primers containing 100 ng Human Genomic DNA with 1 unit of Q5® Hot Start High-Fidelity DNA Polymerase for 25 cycles of PCR amplification results in the expected 665 bp product, and a decrease in non-specific genomic bands after pre-incubation at room temperature for 1 hour, when compared to a non-hot start control reaction.	
PCR Amplification (Enhancer Dependent, >65% GC-rich) A 50 $\mu$ I reaction in Q5® Reaction Buffer and Q5® High GC Enhancer in the presence of 200 $\mu$ M dNTPs and 0.5 $\mu$ M primers containing 20 ng Human Genomic DNA with 1 unit of Q5® Hot Start High-Fidelity DNA Polymerase for 30 cycles of PCR amplification results in the enhancer-dependent production of the expected 452 bp product.	Pass
<b>PCR Amplification (7 kb Human Genomic DNA)</b> A 50 μl reaction in Q5® Reaction Buffer in the presence of 200 μM dNTPs and 0.5 μM primers containing 20 ng Human Genomic DNA with 1 unit of Q5® Hot Start High-Fidelity DNA Polymerase for 30 cycles of PCR amplification results in the expected 7 kb product.	Pass
PCR Amplification (20 kb Lambda DNA) A 50 $\mu$ I reaction in Q5® Reaction Buffer in the presence of 200 $\mu$ M dNTPs and 1.0 $\mu$ M primers containing 10 ng Lambda DNA with 1 unit of Q5® Hot Start High-Fidelity DNA Polymerase for 22 cycles of PCR amplification results in the expected 20 kb product.	Pass
<b>Endonuclease Activity (Hot Start, Nicking)</b> A 50 µl reaction in NEBuffer 2 in the presence of 400 µM dNTPs containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of Q5® High-Fidelity DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>RNase Activity (Extended Digestion)</b> A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of Q5® Hot Start High-Fidelity DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Vayanez hristie

Christie Vazquez Production Scientist 22 Jun 2021

Michae +110

Michael Tonello Packaging Quality Control Inspector 22 Jun 2021

