

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Q5® Hot Start High-Fidelity 2X Master Mix

Catalog Number: M0494L

Concentration: 2 X Concentrate

Packaging Lot Number: 10163755 Expiration Date: 06/2024 Storage Temperature: -20°C

Specification Version: PS-M0494S/L/X v2.0

Composition (1X): Proprietary

Q5® Hot Start High-Fidelity 2X Master Mix Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0494SVIAL	Q5® Hot Start High-Fidelity 2X Master Mix	10156493	Pass	

Assay Name/Specification	Lot # 10163755
PCR Amplification (7 kb Human Genomic DNA, Master Mix) A 50 μl reaction in 1X Q5® Hot Start High-Fidelity Master Mix and 0.5 μM primers containing 20 ng Human Genomic DNA for 30 cycles of PCR amplification results in the expected 7 kb product.	Pass
PCR Amplification (20 kb Lambda DNA, Master Mix) A 50 µl reaction in 1X Q5® Hot Start High-Fidelity Master Mix and 1.0 µM primers containing 10 ng Lambda DNA for 22 cycles of PCR amplification results in the expected 20 kb product.	Pass
Endonuclease Activity (Nicking, Polymerase, dNTP) A 50 μl reaction in NEBuffer 2 in the presence of 400 μM dNTPs containing 1 μg of supercoiled pUC19 DNA and a minimum of 10 units of Q5® High-Fidelity DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 2 units of Q5® High-Fidelity DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass



M0494L / Lot: 10163755

Page 1 of 3

Accov Nama/Specification	L of # 40462755
Assay Name/Specification	Lot # 10163755
Phosphatase Activity (pNPP)	Pass
A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Q5® High-Fidelity DNA	
Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase	
activity as determined by spectrophotometric analysis.	
PCR Amplification (Hot Start, Human Genomic DNA, Master Mix)	Pass
A 25 µl reaction in 1X Q5® Hot Start High-Fidelity Master Mix and 0.5 µM primers	
containing 50 ng Human Genomic DNA for 25 cycles of PCR amplification results in the	
expected 665 bp product and a decrease in non-specific genomic bands after	
pre-incubation at room temperature for 1 hour, when compared to a non-hot start control reaction.	
Some reaction.	
Non-Specific DNase Activity (16 hour, Buffer)	Pass
A 50 µl reaction in 1X Q5® Hot Start High-Fidelity Master Mix containing 1 µg of T3	
or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16	
hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
addition by agained generations.	
Protein Purity Assay (SDS-PAGE)	Pass
Q5® High-Fidelity DNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis	
using Coomassie Blue detection.	
RNase Activity (Extended Digestion)	Pass
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	
and a minimum of 1 µl of Q5® Hot Start High-Fidelity 2X Master Mix is incubated at	
37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as	
determined by gel electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



M0494L / Lot: 10163755

Page 2 of 3

Jan X /2-

Trinh Nguyen Production Scientist 18 Jul 2022 Michael Tonello

Packaging Quality Control Inspector

14 Oct 2022