

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: OneTag® Quick-Load® DNA Polymerase

Catalog Number: M0509S Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 15

nmol of dNTP into acid insoluble material in 30 minutes at 75°C.

Packaging Lot Number: 10156036 Expiration Date: 04/2024 Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.5 % Tween® 20,

0.5 % IGEPAL® CA-630, 50% Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0509S/L/X v2.0

OneTaq® Quick-Load® DNA Polymerase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0509SVIAL	OneTaq® Quick-Load® DNA Polymerase	10147742	Pass	
B9022SVIAL	OneTaq® Standard Reaction Buffer	10143441	Pass	
B0509AVIAL	OneTaq® Quick-Load® Reaction Buffer	10149721	Pass	

Assay Name/Specification	Lot # 10156036
PCR Amplification (5.0 kb Lambda DNA) A 25 μl reaction in OneTaq® Quick-Load® Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 5 ng Lambda DNA with 0.625 units of OneTaq® Quick-Load® DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	Pass
PCR Amplification (5.0 kb Lambda DNA) A 25 μl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 5 ng Lambda DNA with 0.625 units of OneTaq® Quick-Load® DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of OneTaq® Quick-Load® DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass



M0509S / Lot: 10156036

Page 1 of 2

Assay Name/Specification	Lot # 10156036
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a	
reaction containing Lambda-HindIII DNA and a minimum of 5 units of OneTaq®	
Quick-Load® DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern	
free of detectable nuclease degradation as determined by agarose gel	
electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Christie Vazquez Production Scientist 21 Jun 2022

Erin Varney

Packaging Quality Control Inspector

21 Jun 2022

