

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Msz Exonuclease I

Catalog Number: M0527S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will catalyze the

release of 10 nmol of acid-soluble nucleotide in a total reaction volume of 100 μl in 15 minutes at 55°C in 1X rCutSmart Buffer with

0.17 mg/ml single-stranded [3H]-DNA

Packaging Lot Number: 10189662
Expiration Date: 04/2025
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCI, 50 mM KCI, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

(pH 7.4 @ 25°C)

Specification Version: PS-M0527S v2.0

Msz Exonuclease I Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0527SVIAL	Msz Exonuclease I	10183128	Pass	
B6004SVIAL	rCutSmart™ Buffer	10182169	Pass	

Assay Name/Specification	Lot # 10189662
Endonuclease Activity (Circular Single Stranded DNA) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of M13mp18 Single-stranded DNA and a minimum of 100 units of Msz Exonuclease I incubated for 16 hours at 37°C results in <10% conversion to linear DNA as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of Msz Exonuclease I incubated for 16 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Msz Exonuclease I is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	Pass



M0527S / Lot: 10189662 Page 1 of 2

Assay Name/Specification	Lot # 10189662
and a minimum of 1 µl of Msz Exonuclease I is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	
qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of Msz Exonuclease I is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Jenna Ware Production Scientist 24 Apr 2023

genome.

Michael Tonello

Packaging Quality Control Inspector

25 Apr 2023



M0527S / Lot: 10189662

Page 2 of 2