

New England Biolabs Certificate of Analysis

Product Name: *Exonuclease VIII, truncated*
Catalog Number: M0545S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to produce 1 nmol of acid-soluble deoxyribonucleotide from double-stranded DNA in a total reaction volume of 50 µl in 30 minutes at 37°C in 1X NEBuffer 4 with 0.15 mM sonicated duplex [³H]-DNA.
Packaging Lot Number: 10220983
Expiration Date: 03/2026
Storage Temperature: -20°C
Storage Conditions: 50 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.1 % Triton X-100, (pH 7.5 @ 25°C)
Specification Version: PS-M0545S/L v1.0

Exonuclease VIII, truncated Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0545SVIAL	Exonuclease VIII, truncated	10220982	Pass
B7004SVIAL	NEBuffer™ 4	10225682	Pass

Assay Name/Specification	Lot # 10220983
<p>Endonuclease Activity (Circular Single Stranded DNA) A 50 µl reaction in NEBuffer 4 containing 1 µg of M13mp18 Single-stranded DNA and a minimum of 30 units of Exonuclease VIII, truncated incubated for 4 hours at 37°C results in <10% conversion to linear DNA as determined by agarose gel electrophoresis.</p>	Pass
<p>Endonuclease Activity (Nicked Circular DNA) A 50 µl reaction in NEBuffer 4 containing 1 µg of PhiX174 RF II DNA and a minimum of 30 units of Exonuclease VIII, truncated incubated for 4 hours at 37°C results in <10% conversion to linear DNA as determined by agarose gel electrophoresis.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Exonuclease VIII, truncated incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10220983
Protein Purity Assay (SDS-PAGE) Exonuclease VIII, truncated is $\geq 95\%$ pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 10 units of Exonuclease VIII, truncated is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Nicole Castagnozzi
Production Scientist
15 Mar 2024



Michael Tonello
Packaging Quality Control Inspector
15 Mar 2024