

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: EcoGII Methyltransferase

Catalog Number: M0603S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to completely

protect 100 ng of a FAM-labeled dsDNA substrate in 30 minutes at 37°C in a total reaction volume of 20 µL against cleavage by Mbol restriction endonuclease as determined by capillary electrophoresis.

Packaging Lot Number: 10185491
Expiration Date: 03/2025
Storage Temperature: -20°C

Storage Conditions: 250 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

0.15 % TritonX-100, 200 μg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-M0603S/L v1.0

EcoGII Methyltransferase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0603SVIAL	EcoGII Methyltransferase	10183389	Pass	
B9003SVIAL	S-adenosylmethionine (SAM)	10179342	Pass	
B6004SVIAL	rCutSmart™ Buffer	10179340	Pass	

Assay Name/Specification	Lot # 10185491
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled pBR322 DNA and a minimum of 25 units of EcoGII Methyltransferase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 25 units of EcoGII Methyltransferase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (Methyltransferase) A 20 µl reaction in CutSmart® Buffer supplemented with 160 µM SAM containing 100 ng of an 80 bp FAM-labeled dsDNA substrate and 0.5 units of EcoGII Methyltransferase incubated for 30 minutes at 37°C followed by heat inactivation results in ≥ 70%	Pass



M0603S / Lot: 10185491

Page 1 of 2

Assay Name/Specification	Lot # 10185491
protection from digestion with 5 units of Mbol incubated at 37°C for 30 minutes as determined by capillary electrophoresis.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 25 units of EcoGII Methyltransferase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) EcoGII Methyltransferase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of EcoGII Methyltransferase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Mala-Samaranayake-

Production Scientist

16 Mar 2023

Michael Tonello

Packaging Quality Control Inspector

18 Apr 2023



M0603S / Lot: 10185491