

New England Biolabs Certificate of Analysis

Product Name: *EcoGII Methyltransferase*
Catalog Number: M0603S
Concentration: 5,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to completely protect 100 ng of a FAM-labeled dsDNA substrate in 30 minutes at 37°C in a total reaction volume of 20 µL against cleavage by MboI restriction endonuclease as determined by capillary electrophoresis.
Packaging Lot Number: 10263575
Expiration Date: 09/2026
Storage Temperature: -20°C
Storage Conditions: 250 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.15 % Triton X-100, 180 µg/ml BSA, (pH 7.4 @ 25°C)
Specification Version: PS-M0603S/L v3.0


EcoGII Methyltransferase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0603SVIAL	EcoGII Methyltransferase	10253398	Pass
B9003SVIAL	S-adenosylmethionine (SAM)	10233985	Pass
B6004SVIAL	rCutSmart™ Buffer	10250208	Pass

Assay Name/Specification	Lot # 10263575
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 25 units of EcoGII Methyltransferase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 25 units of EcoGII Methyltransferase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 25 units of EcoGII Methyltransferase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel	Pass

Assay Name/Specification	Lot # 10263575
electrophoresis.	
Protein Purity Assay (SDS-PAGE) EcoGII Methyltransferase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of EcoGII Methyltransferase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Jen Wu
Production Scientist
16 Sep 2024



Michael Tonello
Packaging Quality Control Inspector
04 Nov 2024