

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Thermostable FEN1

Catalog Number: M0645S
Concentration: 32,000 U/ml

Unit Definition: One unit is defined as the amount of FEN1 required to cleave 10 pmol

of 5' DNA flap containing oligonucleotide substrate in a total

reaction volume of 10 μl for 10 minutes at 65°C.

Packaging Lot Number: 10098975
Expiration Date: 02/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.1 %

Triton®X-100 , 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0645S/L v1.0

Thermostable FEN1 Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0645SVIAL	Thermostable FEN1	10098976	Pass	
B9004SVIAL	ThermoPol® Reaction Buffer Pack	10099068	Pass	

Assay Name/Specification	Lot # 10098975
Protein Purity Assay (SDS-PAGE) Thermostable FEN1 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie	Pass
Blue detection.	_
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 320 units of Thermostable FEN1 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by gel electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Thermostable FEN1 is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled	Pass



M0645S / Lot: 10098975

Page 1 of 2

Assay Name/Specification	Lot # 10098975
PhiX174 DNA and a minimum of 160 units of Thermostable FEN1 incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 160 units of Thermostable FEN1 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lauren Higgins Production Scientist 10 Mar 2021 Michael Tonello

Packaging Quality Control Inspector

10 Mar 2021



M0645S / Lot: 10098975

Page 2 of 2