

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | EnGen® Spy Cas9 Nickase                                      |
|------------------------|--|
| Catalog Number:        | M0650T   |
| Concentration:         | 20 μΜ  |
| Unit Definition:       | N/A  |
| Packaging Lot Number:  | 10065135   |
| Expiration Date:       | 02/2022  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 % |
|                        | Glycerol, (pH 7.4 @ 25°C)                                    |
| Specification Version: | PS-M0650T v1.0   |

| EnGen® Spy Cas9 Nickase Component List |                         |            |                      |  |
|--|-------------------------|------------|----------------------|--|
| NEB Part Number                        | Component Description   | Lot Number | Individual QC Result |  |
| M0650TVIAL                             | EnGen® Spy Cas9 Nickase | 10065631   | Pass                 |  |
| B7203SVIAL                             | NEBuffer™ 3.1           | 10053975   | Pass                 |  |

| Assay Name/Specification  | Lot # 10065135 |
|---|----------------|
| <b>Functional Testing (EnGen® Spy Cas9 Nickase, Targeted Nicking)</b><br>A 20 µl reaction in 1X NEBuffer 3.1 containing 20 nM 100 bp FAM and ROX labeled<br>double stranded target DNA, 100 nM sgRNA and 100 nM EnGen® Spy Cas9 Nickase<br>incubated for 1 hour at 37°C results in ≥90% nicking of the substrate DNA as<br>determined by capillary electrophoresis. | Pass           |
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1<br>pmol of EnGen® Spy Cas9 Nickase incubated for 16 hours at 37°C results in a DNA<br>pattern free of detectable nuclease degradation as determined by agarose gel<br>electrophoresis.  | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase<br>incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.  | Pass           |
| <b>Endonuclease Activity (Nicking)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 RF I DNA and<br>a minimum of 1 pmol of EnGen® Spy Cas9 Nickase incubated for 4 hours at 37°C results  | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification  | Lot # 10065135 |
|---|----------------|
| in <10% conversion to the nicked form as determined by agarose gel electrophoresis.   |                |
| Protein Purity Assay (SDS-PAGE)<br>EnGen® Spy Cas9 Nickase is ≥ 95% pure as determined by SDS-PAGE analysis using<br>Coomassie Blue detection.  | Pass           |
| <b>RNase Activity (Extended Digestion)</b><br>A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA<br>and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase is incubated at 37°C. After<br>incubation for 16 hours, >90% of the substrate RNA remains intact as determined by<br>gel electrophoresis using fluorescent detection. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

Bhairavi Jani Production Scientist 16 Apr 2020

zay-

Minichiello Packaging Quality Control Inspector 16 Apr 2020

