

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: TelN Protelomerase

Catalog Number: M0651S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 0.5

μg of pMiniT-TeIRL Bsal-linearized DNA in 30 minutes at 30°C in a

total reaction volume of 50 μl.

Packaging Lot Number: 10143307
Expiration Date: 03/2023
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl , 10 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0651S v2.0

TelN Protelomerase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0651SVIAL	TelN Protelomerase	10143306	Pass	
B9004SVIAL	ThermoPol® Reaction Buffer Pack	10139748	Pass	

Assay Name/Specification	Lot # 10143307
Functional Testing (Covalent End Integrity) A 50 μI reaction in ThermoPol® Reaction Buffer containing 0.5 μg of pMiniT-TeIRL DNA and 5 units TeIN Protelomerase incubated for 30 minutes at 30°C followed by heat inactivation and the subsequent addition of 10 units of T5 exonuclease incubated for 1 hour at 37°C results in ≤ 10% loss of starting material as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in ThermoPol® Reaction Buffer containing 1 μg of HaeIII digested PhiX174 RF I DNA and a minimum of 50 units of TeIN Protelomerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in ThermoPol® Reaction Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 25 units of TelN Protelomerase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass



M0651S / Lot: 10143307

Page 1 of 2

Assay Name/Specification	Lot # 10143307
Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of TelN Protelomerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Circular Single Stranded DNA) A 50 μl reaction in ThermoPol® Reaction Buffer containing 1 μg of M13mp18 Single-stranded DNA and a minimum of 25 units of TelN Protelomerase incubated for 4 hours at 37°C results in <20% conversion to linear DNA as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE)  TelN Protelomerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 21 Mar 2022 Michael Tonello

Packaging Quality Control Inspector

21 Mar 2022

M0651S / Lot: 10143307