

DVERY 240 County Road INE Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Faustovirus Capping Enzyme

Catalog Number: M2081S
Concentration: 25,000 U/ml

Unit Definition: One unit of Faustovirus Capping Enzyme is defined as the amount of

enzyme required to convert 75 pmol of a 20-mer transcript to Cap-0

RNA in 30 minutes at 37°C.

Packaging Lot Number: 10185135
Expiration Date: 02/2025
Storage Temperature: -20°C

Storage Conditions: 40 mM Tris-HCl, 100 mM NaCl, 50 mM Arginine, 0.1 mM TCEP, 50%

Glycerol, (pH 8.0 @ 25°C)

Specification Version: PS-M2081S/L v1.0

| Faustovirus Capping Enzyme Component List |                            |            |                      |  |
|---|----------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>                    | Component Description      | Lot Number | Individual QC Result |  |
| N2080AVIAL                                | GTP                        | 10176039   | Pass                 |  |
| M2081SVIAL                                | Faustovirus Capping Enzyme | 10178541   | Pass                 |  |
| B9003SVIAL                                | S-adenosylmethionine (SAM) | 10171171   | Pass                 |  |
| B2181AVIAL                                | FCE Capping Buffer         | 10181122   | Pass                 |  |

| Assay Name/Specification  | Lot # 10185135 |
|---|----------------|
| Endonuclease Activity (Nicking) A 50 μl reaction in FCE Capping Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 25 units of Faustovirus Capping Enzyme incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass           |
| Endonuclease Activity (Nicking) A 50 μl reaction in FCE Capping Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 25 units of Faustovirus Capping Enzyme incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in FCE Capping Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 25 units of Faustovirus Capping Enzyme incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.   | Pass           |



M2081S / Lot: 10185135

Page 1 of 3

| Assay Name/Specification   | Lot # 10185135 |
|--|----------------|
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in FCE Capping Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 25 units of Faustovirus Capping Enzyme incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.  | Pass           |
| Protein Purity Assay (SDS-PAGE) Faustovirus Capping Enzyme is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.  | Pass           |
| Protein Purity Assay (SDS-PAGE) Faustovirus Capping Enzyme is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.  | Pass           |
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 25 units of Faustovirus Capping Enzyme is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.   | Pass           |
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 25 units of Faustovirus Capping Enzyme is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.   | Pass           |
| <b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 25 units of Faustovirus Capping Enzyme is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome. | Pass           |
| <b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 25 units of Faustovirus Capping Enzyme is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit



M2081S / Lot: 10185135 Page 2 of 3



www.neb.com/trademarks for additional information.

Jessica Cane Production Scientist

01 Mar 2023

Michael Tonello

Packaging Quality Control Inspector

22 Mar 2023

Page 3 of 3