

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Hi-T4™ DNA Ligase
Catalog Number:	M2622S
Concentration:	400,000 U/mI
Unit Definition:	One unit is defined as the amount of enzyme required to give 50% ligation of 6 μg of Lambda-HindIII DNA in 30 minutes at 25°C in a total reaction volume of 20 μl.
Packaging Lot Number:	10092876
Expiration Date:	12/2022
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version:	PS-M2622S/L v1.0

Hi-T4 [™] DNA Ligase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M2622SVIAL	Hi-T4™ DNA Ligase	10092875	Pass	
B0535AVIAL	StickTogether™ DNA Ligase Buffer	10077408	Pass	
B0202AVIAL	T4 DNA Ligase Reaction Buffer	10086512	Pass	

Assay Name/Specification	Lot # 10092876
DNase Activity (Labeled Oligo, 5' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 2000 units of Hi-T4 [™] DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
Double Stranded DNase Activity (Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 2000 units of Hi-T4 [™] DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Hi-T4 [™] DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel	Pass





be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10092876
electrophoresis using fluorescent detection.	
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 400 units of Hi-T4™ DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
DNase Activity (Labeled Oligo, 3' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 2000 units of Hi-T4 [™] DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 2000 units of Hi-T4 [™] DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
Protein Concentration (A280) The concentration of Hi-T4 [™] DNA Ligase is 0.4 mg/ml +/- 10% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 57,675 and molecular weight of 56,806 daltons for Hi-T4 [™] DNA Ligase (Pace, C.N. et al. (1995) Protein Sci., 4, 2411-2423).	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1 containing 1 µg of CIP-treated Lambda-HindIII DNA and a minimum of 400 units of Hi-T4 [™] DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Hi-T4 [™] DNA Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 400 units of Hi-T4 [™] DNA Ligase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.





be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 11 Jan 2021

Ton. N. Michae

Michael Tonello Packaging Quality Control Inspector 11 Jan 2021

