

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Thermolabile USER® II Enzyme
Catalog Number:	M5508S
Concentration:	1,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to nick 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base in 15 minutes at 37°C in a total reaction volume of 10 μL in 1X T4 DNA Ligase Buffer.
Packaging Lot Number:	10066485
Expiration Date:	08/2021
Storage Temperature:	-20°C
Storage Conditions:	25 mM KCl, 35 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 15 mM Tris-HCl, 100 μg/ml BSA, 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version:	PS-M5508S/L v1.0

Thermolabile USER® II Enzyme Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M5508SVIAL	Thermolabile USER® II Enzyme	10054426	Pass	
B7204SVIAL	CutSmart® Buffer	10061304	Pass	

Assay Name/Specification	Lot # 10066485
<b>Functional Testing (Thermolability, Endonuclease III)</b> A 10 µl reaction in CutSmart® Buffer containing 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base and 1 unit of Thermolabile USER® II Enzyme was incubated for 15 minutes at 37°C followed by heat inactivation for 10 minutes at 65°C. The addition of 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single AP site and incubation for 15 minutes at 37°C followed by 10 minutes at 75°C, results in no cleavage of additional substrate.	Pass
<b>Functional Testing (Thermolability, UDG)</b> A 10 µl reaction in CutSmart® Buffer containing 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base and 1 unit of Thermolabile USER® II Enzyme was incubated for 15 minutes at 37°C followed by heat inactivation for 10 minutes at 65°C. The addition of 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base with 20 units of Endonuclease III and incubation for 15 minutes at 37°C followed by 10 minutes at 75°C, results in no cleavage of additional substrate.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10066485
<b>Functional Testing (USER, Transformation assay)</b> A 10 µl reaction in ThermoPol® Reaction Buffer containing 20 ng linearized pNEB206A, 100 ng of a 950 bp control PCR product and 1 unit of Thermolabile USER® II Enzyme was incubated for 15 minutes at 37°C followed by 15 minutes at 25°C. After transformation into ER2267 chemically-competent cells >95% of colonies contained recombinant plasmid.	Pass
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 1 unit of Thermolabile USER® II Enzyme is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is $\leq$ 1 E. coli genome.	Pass
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Thermolabile USER® II Enzyme is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

dauren biggins

Lauren Sears Higgins Production Scientist 30 Aug 2019

Michae MI.

Michael Tonello Packaging Quality Control Inspector 27 Jan 2020

