

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Thermolabile USER® II Enzyme
Catalog Number:	M5508S
Concentration:	1,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to nick 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base in 15 minutes at 37°C in a total reaction volume of 10 μL in 1X T4 DNA Ligase Buffer.
Packaging Lot Number:	10182213
Expiration Date:	02/2025
Storage Temperature:	-20°C
Storage Conditions:	25 mM KCl, 35 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 15 mM Tris-HCl, 100 μg/ml BSA, 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version:	PS-M5508S/L v1.0

Thermolabile USER® II Enzyme Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M5508SVIAL	Thermolabile USER® II Enzyme	10178402	Pass	
B6004SVIAL	rCutSmart™ Buffer	10175292	Pass	

Assay Name/Specification	Lot # 10182213
Functional Testing (USER, Transformation assay) A 10 µl reaction in ThermoPol® Reaction Buffer containing 20 ng linearized pNEB206A, 100 ng of a 950 bp control PCR product and 1 unit of Thermolabile USER® II Enzyme was incubated for 15 minutes at 37°C followed by 15 minutes at 25°C. After transformation into ER2267 chemically-competent cells >95% of colonies contained recombinant plasmid.	Pass
Functional Testing (Thermolability, UDG) A 10 µl reaction in CutSmart® Buffer containing 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base and 1 unit of Thermolabile USER® II Enzyme was incubated for 15 minutes at 37°C followed by heat inactivation for 10 minutes at 65°C. The addition of 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base with 20 units of Endonuclease III and incubation for 15 minutes at 37°C followed by 10 minutes at 75°C, results in no cleavage of additional substrate.	Pass
Functional Testing (Thermolability, Endonuclease III)	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10182213
A 10 µl reaction in CutSmart® Buffer containing 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base and 1 unit of Thermolabile USER® II Enzyme was incubated for 15 minutes at 37°C followed by heat inactivation for 10 minutes at 65°C. The addition of 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single AP site and incubation for 15 minutes at 37°C followed by 10 minutes at 75°C, results in no cleavage of additional substrate.	
RNase Activity (Extended Digestion) A 10 μ I reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ I of Thermolabile USER® II Enzyme is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 1 unit of Thermolabile USER® II Enzyme is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is \leq 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lauren Higgins

Lauren Sears Higgins Production Scientist 24 Feb 2023

Minhae

Michael Tonello Packaging Quality Control Inspector 27 Feb 2023

