

New England Biolabs Certificate of Analysis

Product Name: M-MuLV Reverse Transcriptase
Catalog Number: M0253S
Concentration: 200,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to incorporate 1 nmol of dTTP into an acid-insoluble form in 10 minutes at 37°C.
Packaging Lot Number: 10135007
Expiration Date: 12/2023
Storage Temperature: -20°C
Storage Conditions: 50 mM Tris-HCl, 150 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 0.1 % IGEPAL® CA-630, 50 % Glycerol, (pH 7.6 @ 25°C)
Specification Version: PS-M0253S/L v2.0

M-MuLV Reverse Transcriptase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0253SVIAL	M-MuLV Reverse Transcriptase	10135565	Pass
B0253SVIAL	M-MuLV Reverse Transcriptase Reaction Buffer	10131962	Pass

Assay Name/Specification	Lot # 10135007
<p>RNase Activity Assay (2 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of M-MuLV Reverse Transcriptase incubated for 2 hours at 37°C results in no detectable degradation of the RNA as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in M-MuLV Reverse Transcriptase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of M-MuLV Reverse Transcriptase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in M-MuLV Reverse Transcriptase Reaction Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 200 units of M-MuLV Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10135007
<p>Endonuclease Activity (Nicking) A 50 µl reaction in M-MuLV Reverse Transcriptase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 200 units of M-MuLV Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Christie Vazquez
Production Scientist
24 Feb 2022



Josh Hersey
Packaging Quality Control Inspector
24 Feb 2022