

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Antarctic Thermolabile UDG

Catalog Number: M0372S Concentration: 1,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the

release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [ 3H] -uracil in a 50 μl reaction containing 0.2 μg DNA (104-105 cpm/μg)

in 30 minutes at 37°C.

Packaging Lot Number: 10084390 Expiration Date: 09/2022 Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

(pH 7.4 @ 25°C)

Specification Version: PS-M0372S/L v1.0

| Antarctic Thermolabile UDG Component List |                                   |            |                      |  |
|---|-----------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>                    | Component Description             | Lot Number | Individual QC Result |  |
| M0372SVIAL                                | Antarctic Thermolabile UDG        | 10084391   | Pass                 |  |
| B9014SVIAL                                | Standard Taq Reaction Buffer Pack | 10073672   | Pass                 |  |

| Assay Name/Specification   | Lot # 10084390 |
|--|----------------|
| DNase Activity (Labeled Oligo, 3' extension) A 50 μl reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 1 unit of Antarctic Thermolabile UDG incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis. | Pass           |
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of f-300 RNA transcript and a minimum of 1 unit of Antarctic Thermolabile UDG is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using agarose gel electrophoresis.             | Pass           |
| Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 1 unit of Antarctic Thermolabile UDG incubated for 16 hours at 37°C yields <5% degradation as determined by capillary   | Pass           |



M0372S / Lot: 10084390

Page 1 of 3

| Assay Name/Specification   | Lot # 10084390 |
|--|----------------|
| electrophoresis.   |                |
| DNase Activity (Labeled Oligo, 5' extension) A 50 µl reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 1 unit of Antarctic Thermolabile UDG incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.   | Pass           |
| Double Stranded DNase Activity (Labeled Oligo) A 50 µl reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 1 unit of Antarctic Thermolabile UDG incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.  | Pass           |
| Endonuclease Activity (Nicking) A 50 µl reaction in Standard Taq Reaction Buffer containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 15 units of Antarctic Thermolabile UDG incubated for 4 hours at 37°C results in <20% conversion to RFII as determined by agarose gel electrophoresis.  | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in Standard Taq Reaction Buffer containing 1 µg of HindIII digested Lambda DNA and a minimum of 50 units of Antarctic Thermolabile UDG incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.  | Pass           |
| Protein Purity Assay (SDS-PAGE) Antarctic Thermolabile UDG is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.  | Pass           |
| <b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 1 unit of Antarctic Thermolabile UDG is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



M0372S / Lot: 10084390

Lauren Higgins

Lauren Higgins Production Scientist 23 Sep 2020 Michael Tonello

Packaging Quality Control Inspector

23 Sep 2020