

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: beta-Agarase I

Catalog Number: M0392S Concentration: 1,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 200

μl of molten low melting point or NuSieve agarose to nonprecipitable

neoagaro-oligosaccharides in 1 hour at 42°C

Packaging Lot Number: 10191395 Expiration Date: 05/2025 Storage Temperature: -20°C

Storage Conditions: 50 mM Bis-Tris-HCl, 1 mM EDTA, 50 % Glycerol, (pH 6.5 @ 25°C)

Specification Version: PS-M0392S/L v1.0

beta-Agarase I Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0392SVIAL	β-Agarase I	10188114	Pass	
B0392SVIAL	ß-Agarase I Reaction Buffer	10188841	Pass	

Assay Name/Specification	Lot # 10191395
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and	Pass
a minimum of 1 unit of β-Agarase I incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 5 units of β-Agarase I incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda DNA and a minimum of 10 units of β-Agarase I incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) β-Agarase I is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



M0392S / Lot: 10191395

Page 1 of 2

Assay Name/Specification	Lot # 10191395
RNase Activity (Extended Digestion)	Pass
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	
and a minimum of 1 μl of β-Agarase I is incubated at 37°C. After incubation for 16	
hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis	
using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Vanessa Mathieu-Sheltry Production Scientist 04 May 2023

Mulhiu hr

Josh Hersey

Packaging Quality Control Inspector

19 May 2023

