

New England Biolabs Certificate of Analysis

Product Name: 7-deaza-dGTP
Catalog Number: N0445S
Concentration: 5 mM
Unit Definition: N/A
Packaging Lot Number: 10244073
Expiration Date: 02/2026
Storage Temperature: -20°C
Storage Conditions: Supplied in Ultrapure water as a lithium salt , (pH 7.0)
Specification Version: PS-N0445S/L v2.0


7-deaza-dGTP Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N0445SVIAL	7-deaza-dGTP	10226743	Pass

Assay Name/Specification	Lot # 10244073
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 µl of 7-deaza-dGTP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 µl of 7-deaza-dGTP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (0.5 kb Lambda DNA, 7-deaza) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 µM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (2 kb Lambda DNA, 7-deaza) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 µM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 2 kb product.	Pass

Assay Name/Specification	Lot # 10244073
<p>PCR Amplification (5 kb Lambda DNA, 7-deaza) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 µM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.</p>	Pass
<p>Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 80 µl 7-deaza-dGTP incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>Physical Purity (HPLC) 7-deaza-dGTP is ≥ 95% pure as determined by HPLC analysis.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of 7-deaza-dGTP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.


Lea Antonopoulos
Production Scientist
13 Mar 2024


Michael Tonello
Packaging Quality Control Inspector
03 Jun 2024