

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Deoxynucleotide (dNTP) Solution Set
Catalog #:	N0446S
Concentration:	100 mM
Unit Definition:	N/A
<i>Lot</i> #:	0551602
Assay Date:	02/2016
Expiration Date:	02/2018
Storage Temp:	-20°C
Storage Conditions:	Supplied in Ultrapure water as a sodium salt (pH 7.5)
Specification Version:	PS-N0446S v1.0
Effective Date:	23 Feb 2016

Assay Name/Specification (minimum release criteria)	Lot #0551602
Endonuclease Activity (Nicking) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 1 μ l of dATP, dCTP, dGTP, and dTTP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 μ l of dATP, dCTP, dGTP, and dTTP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (0.5 kb Lambda, dNTPs) - A 50 μ l reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M dATP, dCTP, dGTP, and dTTP and 0.5 μ M primers containing 1 ng Lambda DNA with 1.25 units of <i>Taq</i> DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (2.0 kb Lambda, dNTPs) - A 50 μ l reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M dATP, dCTP, dGTP, and dTTP and 0.5 μ M primers containing 1 ng Lambda DNA with 1.25 units of <i>Taq</i> DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product.	Pass
PCR Amplification (5.0 kb Lambda, dNTPs) - A 50 μ l reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M dATP, dCTP, dGTP, and dTTP and 0.5 μ M primers containing 1 ng Lambda DNA with 1.25 units of <i>Taq</i> DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	Pass
Phosphatase Activity (pNPP) - A 200 μ l reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 mM <i>p</i> -Nitrophenyl Phosphate (pNPP) and a minimum of 4 μ l dATP, dCTP, dGTP, and dTTP incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass



N0446S Lot: 0551602 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0551602
Physical Purity (HPLC) - dATP, dCTP, dGTP, and dTTP is \geq 99% pure as determined by HPLC analysis.	Pass
RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single- stranded RNA and a minimum of 1 μ l dATP, dCTP, dGTP, and dTTP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

N

Authorized by Melanie Fortier 23 Feb 2016



Inspected by Denisa Gilaj 12 Jul 2016