

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Deoxynucleotide (dNTP) Solution Set

Catalog Number: N0446S
Concentration: 100 mM
Unit Definition: N/A

Packaging Lot Number: 10122833
Expiration Date: 03/2023
Storage Temperature: -20°C

Storage Conditions: Supplied in Ultrapure water as a sodium salt (pH 7.5)

Specification Version: PS-N0446S v2.0

Deoxynucleotide (dNTP) Solution Set Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
N0443SVIAL	dTTP	10105791	Pass	
N0442SVIAL	dGTP	10105792	Pass	
N0441SVIAL	dCTP	10105788	Pass	
N0440SVIAL	dATP Solution	10105793	Pass	

Assay Name/Specification	Lot # 10122833
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 4 µl dATP, dCTP, dGTP, and dTTP incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Physical Purity (HPLC) dATP, dCTP, dGTP, and dTTP is ≥ 99% pure as determined by HPLC analysis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl dATP, dCTP, dGTP, and dTTP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
PCR Amplification (2.0 kb Lambda, dNTPs) A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dGTP, and dTTP and 0.5 μM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb	Pass



N0446S / Lot: 10122833

Page 1 of 2

Assay Name/Specification	Lot # 10122833
product.	
PCR Amplification (0.5 kb Lambda, dNTPs) A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dGTP, and dTTP and 0.5 μM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (5.0 kb Lambda, dNTPs) A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dGTP, and dTTP and 0.5 μM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 1 μl of dATP, dCTP, dGTP, and dTTP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 µl of dATP, dCTP, dGTP, and dTTP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Christie Vazquez **Production Scientist** 04 Oct 2021

haistie Vazguez

Michael Tonello

Packaging Quality Control Inspector

04 Oct 2021



N0446S / Lot: 10122833

Page 2 of 2