

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Deoxynucleotide (dNTP) Solution Mix
Catalog Number:	N0447L
Concentration:	10 mM
Unit Definition:	N/A
Packaging Lot Number:	10086435
Expiration Date:	09/2022
Storage Temperature:	-20°C
Storage Conditions:	Supplied in Ultrapure water as a sodium salt (pH 7.5)
Specification Version:	PS-N0447S/L v3.0

Deoxynucleotide (dNTP) Solution Mix Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
N0447LVIAL	Deoxynucleotide (dNTP) Solution Mix	10083859	Pass	

Assay Name/Specification	Lot # 10086435
Phosphatase Activity (pNPP) A 200 μl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 μl Deoxynucleotide (dNTP) Solution Mix incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Physical Purity (HPLC) Deoxynucleotide (dNTP) Solution Mix is \geq 99% pure as determined by HPLC analysis.	Pass
RNase Activity (Extended Digestion) A 10 μ I reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ I of Deoxynucleotide (dNTP) Solution Mix is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Endonuclease Activity (Nicking) A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 10 μ l of Deoxynucleotide (dNTP) Solution Mix incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour)	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10086435
A 50 μ I reaction in NEBuffer 2 containing 1 μ g of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 μ I of Deoxynucleotide (dNTP) Solution Mix incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
PCR Amplification (0.5 kb Lambda, dNTPs) A 50 μ l reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M Deoxynucleotide (dNTP) Solution Mix and 0.5 μ M primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (2.0 kb Lambda, dNTPs) A 50 μ l reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M Deoxynucleotide (dNTP) Solution Mix and 0.5 μ M primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product.	Pass
PCR Amplification (5.0 kb Lambda, dNTPs) A 50 μ I reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M Deoxynucleotide (dNTP) Solution Mix and 0.5 μ M primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

pristie Vayanez_

Christie Vazquez Production Scientist 13 Oct 2020

Josh Hersey

Packaging Quality Control Inspector 13 Oct 2020



N0447L / Lot: 10086435 Page 2 of 2