

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

HeLa Genomic DNA
N4006S
100 µg/ml
N/A
0051711
11/2017
11/2019
-20°C
10 mM Tris-HCl , 1 mM EDTA, (pH 7.5 @ 25°C)
PS-N4006S v1.0
07 Jan 2016

Assay Name/Specification (minimum release criteria)	Lot #0051711
<b>A260/A280 Assay</b> - The ratio of UV absorption of HeLa Genomic DNA at 260 and 280 nm is between 1.8 and 2.0.	Pass
<b>DNA Concentration (A260)</b> - The concentration of HeLa Genomic DNA is between 100 and 110 $\mu$ g/ml as determined by UV absorption at 260 nm.	Pass
<b>Electrophoretic Pattern (Genomic DNA)</b> - The banding pattern of HeLa Genomic DNA on a 1.2% agarose gel is evaluated against a control lot for relative integrity and intensity as determined by gel electrophoresis using Ethidium Bromide.	Pass
<b>Non-Specific DNase Activity (Genomic DNA, 16 hour)</b> - A 50 $\mu$ l reaction in 1X NEBuffer 2 containing 2.5 $\mu$ g of HeLa Genomic DNA incubated for 16 hours at 37°C does not produce any further detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Restriction Digest (Genomic DNA)</b> - A 50 µl reaction in NEBuffer 2.1 containing 2.5 µg of HeLa Genomic DNA and 20 units of HindIII incubated for 1 hour at 37°C produces the expected fragmentation pattern as determined by agarose gel electrophoresis.	Pass

otunon

Authorized by Derek Robinson 07 Jan 2016



Mulhicehite

Inspected by Vanessa Mathieu-Sheltry 07 Nov 2017