

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	M13mp18 Single-stranded DNA
Catalog Number:	N4040S
Concentration:	250 μg/ml
Unit Definition:	N/A
Lot Number:	10052522
Expiration Date:	04/2021
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCI (pH 8.0), 1 mM EDTA
Specification Version:	PS-N4040S v2.0

M13mp18 Single-stranded DNA Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
N4040SVIAL	M13mp18 Single-stranded DNA	10042807	Pass	

Assay Name/Specification	Lot # 10052522
<b>Non-Specific DNase Activity (DNA, 16 hour)</b> A 50 µl reaction in 1X NEBuffer 2 containing 2.5 µg of M13mp18 Single-stranded DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Restriction Digest (Single Stranded, Resistant) A 50 µl reaction in CutSmart™ Buffer containing 2.5 µg of M13mp18 Single-stranded DNA and a minimum of 20 units of Xhol incubated for 1 hour at 37°C results in no detectable digestion of the DNA as determined by agarose gel electrophoresis.	Pass
<b>A260/A280 Assay</b> The ratio of UV absorption of M13mp18 Single-stranded DNA at 260 and 280 nm is between 1.8 and 2.0.	Pass
DNA Concentration (A260) The concentration of M13mp18 Single-stranded DNA is between 250 and 260 µg/ml as determined by UV absorption at 260 nm.	Pass
<b>Electrophoretic Pattern (Plasmid)</b> The banding pattern of M13mp18 Single-stranded DNA on a 1.2% agarose gel is evaluated against a control lot for sharpness and relative intensity as determined by gel electrophoresis using Ethidium Bromide.	Pass





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10052522
<b>Mung Bean Nuclease Digest (Sensitive)</b> A 100 µl reaction in Mung Bean Nuclease Reaction Buffer containing 2.5 µg of M13mp18 Single-stranded DNA and 10 units of Mung Bean Nuclease incubated for 1 hour at 30°C results in complete digestion of the DNA as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Nulhice

Vanessa Mathieu-Sheltry Production Scientist 18 Apr 2019

 $a_{l}$ 

Minichiello Packaging Quality Control Inspector 19 Aug 2019

