

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: M13mp18 Single-stranded DNA

Catalog Number:N4040SConcentration:250 μg/ml

Unit Definition: N/A

Packaging Lot Number: 10128627
Expiration Date: 11/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl (pH 8.0), 1 mM EDTA

Specification Version: PS-N4040S v2.0

| M13mp18 Single-stranded DNA Component List | | | | |
|--|-----------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| N4040SVIAL | M13mp18 Single-stranded DNA | 10128626 | Pass | |

| Assay Name/Specification | Lot # 10128627 |
|--|----------------|
| A260/A280 Assay The ratio of UV absorption of M13mp18 Single-stranded DNA at 260 and 280 nm is between 1.8 and 2.0. | Pass |
| DNA Concentration (A260) The concentration of M13mp18 Single-stranded DNA is between 250 and 260 μg/ml as determined by UV absorption at 260 nm. | Pass |
| Electrophoretic Pattern (Plasmid) The banding pattern of M13mp18 Single-stranded DNA on a 1.2% agarose gel is evaluated against a control lot for sharpness and relative intensity as determined by gel electrophoresis using Ethidium Bromide. | Pass |
| Non-Specific DNase Activity (DNA, 16 hour) A 50 µl reaction in 1X NEBuffer 2 containing 2.5 µg of M13mp18 Single-stranded DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Mung Bean Nuclease Digest (Sensitive) A 100 μl reaction in Mung Bean Nuclease Reaction Buffer containing 2.5 μg of M13mp18 Single-stranded DNA and 10 units of Mung Bean Nuclease incubated for 1 hour at 30°C results in complete digestion of the DNA as determined by agarose gel | Pass |



N4040S / Lot: 10128627

Page 1 of 2

| Assay Name/Specification | Lot # 10128627 |
|--|----------------|
| electrophoresis. | |
| Restriction Digest (Single Stranded, Resistant) | Pass |
| A 50 µl reaction in CutSmart™ Buffer containing 2.5 µg of M13mp18 Single-stranded DNA and a minimum of 20 units of Xhol incubated for 1 hour at 37°C results in no | |
| detectable digestion of the DNA as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Vanessa Mathieu-Sheltry
Production Scientist

22 Nov 2021

Michael Tonello

Packaging Quality Control Inspector

22 Nov 2021