

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	PNGase A
Catalog Number:	P0707L
Concentration:	5,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 1 μ g of denatured recombinant Avidin produced in Maize in 1 hour at 37°C in a total reaction volume of 10 μ l.
Lot Number:	10025488
Expiration Date:	10/2019
Storage Temperature:	4°C
Storage Conditions:	50 mM NaCl, 20 mM Tris-HCl, 5 mM EDTA, (pH 7.5 @ 25°C)
Specification Version:	PS-P0707S/L v1.0

PNGase A Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
P0707LVIAL	PNGase A	10023226	Pass	
B2704SVIAL	NP-40	10021820	Pass	
B1720SVIAL	10X Glyco Buffer 3	10022682	Pass	
B1704SVIAL	Glycoprotein Denaturing Buffer	10017111	Pass	

Assay Name/Specification	Lot # 10025488
Glycosidase Activity (α-Glucosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Glucosidase substrate (Glc α 1-6Glc α 1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-Neuraminidase) A 10 μ l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Neuraminidase substrate (Neu5Ac α 2-3Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-2 Fucosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-2Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer	Pass





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10025488
chromatography.	
Glycosidase Activity (α1-3 Fucosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Galactosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-3Gal β 1-4GlcNAc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Mannosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-3Man β 1-4GlcNAc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-6 Galactosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-6Gal α 1-6Glc α 1-2Fru-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-6 Mannosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α -N-Acetylgalactosaminidase substrate (GalNAc α 1-3(Fuc α 1-2)Gal β 1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Mannosidase) A 10 μ I reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β -Mannosidase substrate (Man β 1-4Man β 1-4Man-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10025488
Glycosidase Activity (β-Xylosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β -Xylosidase substrate (Xyl β 1-4Xyl β 1-4Xyl β 1-4Xyl-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-3 Galactosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β -Galactosidase substrate (Gal β 1-3GlcNAc β 1-4Gal β 1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μ I reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β -N-Acetylglucosaminidase substrate (GlcNAc β 1-4GlcNAc β 1-4GlcNAc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Protease Activity (SDS-PAGE) A 20 µl reaction in 1X Glyco Buffer 3 containing 24 µg of a standard mixture of proteins and a minimum of 25 units of PNGase A incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Protein Purity Assay (SDS-PAGE) PNGase A is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

grd

Brad Landgraf Production Scientist 31 Jul 2018

Michae 11.

Michael Tonello Packaging Quality Control Inspector 22 Jan 2019

