

## New England Biolabs Certificate of Analysis

**Product Name:** PNGase A  
**Catalog Number:** P0707S  
**Concentration:** 5,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 1 µg of denatured recombinant Avidin produced in Maize in 1 hour at 37°C in a total reaction volume of 10 µl.  
**Packaging Lot Number:** 10190540  
**Expiration Date:** 04/2024  
**Storage Temperature:** 4°C  
**Storage Conditions:** 50 mM NaCl, 20 mM Tris-HCl, 5 mM EDTA, (pH 7.5 @ 25°C)  
**Specification Version:** PS-P0707S/L v1.0

| PNGase A Component List |                                |            |                      |
|-------------------------|--------------------------------|------------|----------------------|
| NEB Part Number         | Component Description          | Lot Number | Individual QC Result |
| P0707SVIAL              | PNGase A                       | 10184070   | Pass                 |
| B2704SVIAL              | NP-40                          | 10161534   | Pass                 |
| B1720SVIAL              | 10X Glycobuffer 3              | 10120516   | Pass                 |
| B1704SVIAL              | Glycoprotein Denaturing Buffer | 10181130   | Pass                 |


| Assay Name/Specification   | Lot # 10190540 |
|--|----------------|
| <b>Glycosidase Activity (α-Glucosidase)</b><br>A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.   | Pass           |
| <b>Glycosidase Activity (α-N-Acetylgalactosaminidase)</b><br>A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fucα1-2)Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass           |
| <b>Glycosidase Activity (α-Neuraminidase)</b><br>A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as  | Pass           |

| Assay Name/Specification   | Lot # 10190540 |
|--|----------------|
| determined by thin layer chromatography.   |                |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-2 Fucosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuca1-2Gal<math>\beta</math>1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>  | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-3 Fucosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuca1-3Gal<math>\beta</math>1-4GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-3 Galactosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-3Gal<math>\beta</math>1-4GlcNAc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>                         | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-3 Mannosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-3Man<math>\beta</math>1-4GlcNAc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>                             | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-6 Galactosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-6Gal<math>\alpha</math>1-6Glc<math>\alpha</math>1-2Fru-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>  | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\alpha</math>1-6 Mannosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-6Man<math>\alpha</math>1-6(Man<math>\alpha</math>1-3)Man-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>    | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\beta</math>-Mannosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\beta</math>-Mannosidase substrate (Man<math>\beta</math>1-4Man<math>\beta</math>1-4Man-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>                                      | <b>Pass</b>    |

| Assay Name/Specification  | Lot # 10190540 |
|---|----------------|
| <p><b>Glycosidase Activity (<math>\beta</math>-N-Acetylgalactosaminidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\beta</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>    | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\beta</math>-N-Acetylglucosaminidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\beta</math>-N-Acetylglucosaminidase substrate (GlcNAc<math>\beta</math>1-4GlcNAc<math>\beta</math>1-4GlcNAc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>  | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\beta</math>-Xylosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\beta</math>-Xylosidase substrate (Xyl<math>\beta</math>1-4Xyl<math>\beta</math>1-4Xyl<math>\beta</math>1-4Xyl-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>             | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\beta</math>1-3 Galactosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | <b>Pass</b>    |
| <p><b>Glycosidase Activity (<math>\beta</math>1-4 Galactosidase)</b><br/>A 10 <math>\mu</math>l reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-4GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 10 units of PNGase A incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | <b>Pass</b>    |
| <p><b>Protease Activity (SDS-PAGE)</b><br/>A 20 <math>\mu</math>l reaction in 1X Glyco Buffer 3 containing 24 <math>\mu</math>g of a standard mixture of proteins and a minimum of 25 units of PNGase A incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>  | <b>Pass</b>    |
| <p><b>Protein Purity Assay (SDS-PAGE)</b><br/>PNGase A is <math>\geq</math> 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>   | <b>Pass</b>    |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



---

Maxwell Elkus  
Production Scientist  
19 Apr 2023



---

Michael Tonello  
Packaging Quality Control Inspector  
06 Jul 2023