

New England Biolabs Certificate of Analysis

Product Name: Rapid™ PNGase F (non-reducing format)
Catalog Number: P0711S
Unit Definition: N/A
Packaging Lot Number: 10075418
Expiration Date: 06/2021
Storage Temperature: 4°C
Storage Conditions: 50 mM NaCl , 20 mM Tris-HCl , 5 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0711S v2.0

Rapid™ PNGase F (non-reducing format) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0711SVIAL	Rapid™ PNGase F (non-reducing format)	10075417	Pass
B0717SVIAL	5X Rapid PNGase F Buffer (non-reducing format)	10076469	Pass

Assay Name/Specification	Lot # 10075418
Glycosidase Activity (Endo F1, F2, H) A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-3 Galactosidase) A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Xylosidase) A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 1 µl of Rapid PNGase F (non-reducing format)	Pass

Assay Name/Specification	Lot # 10075418
<p>incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Mannosidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Mannosidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Galactosidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Mannosidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Galactosidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Fucosidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled α-Fucosidase substrate</p>	Pass

Assay Name/Specification	Lot # 10075418
<p>(Fuca1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (α1-2 Fucosidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-2Galβ1-4Glc-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Neuraminidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fuca1-2)Galβ1-4Glc-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Glucosidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (Endo F2, F3) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) Rapid PNGase F (non-reducing format) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Protease Activity (SDS-PAGE) A 20 μl reaction in 1X Rapid PNGase F Buffer (non-reducing format) containing 24 μg of a standard mixture of proteins and a minimum of 5 μl of Rapid PNGase F</p>	Pass

Assay Name/Specification	Lot # 10075418
<p>(non-reducing format) incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p> <p>Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 1 μl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.



Alicia Bielik
Production Scientist
30 Jun 2020



Michael Tonello
Packaging Quality Control Inspector
30 Jun 2020