

New England Biolabs Certificate of Analysis

Product Name: β 1-3 Galactosidase
Catalog Number: P0726S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to cleave > 95% of the terminal β -D-galactose from 1 nmol of Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μ l.
Packaging Lot Number: 10084147
Expiration Date: 09/2022
Storage Temperature: -20°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0726S/L v1.0

β1-3 Galactosidase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0726SVIAL	β 1-3 Galactosidase	10084148	Pass
B9001SVIAL	Purified BSA	10059427	Pass
B1727SVIAL	10X GlycoBuffer 1	10066905	Pass

Assay Name/Specification	Lot # 10084147
Protein Purity Assay (SDS-PAGE) β 1-3 Galactosidase is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Protease Activity (SDS-PAGE) A 20 μ l reaction in 1X Glyco Buffer 1 containing 24 μ g of a standard mixture of proteins and a minimum of 100 units of β 1-3 Galactosidase incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Glycosidase Activity (α1-3 Fucosidase) A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 100 units of β 1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (Endo F1, F2, H)	Pass

Assay Name/Specification	Lot # 10084147
<p>A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (Endo F2, F3) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (PNGase F) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Glucosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fuca1-2)Galβ1-4Glc-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Neuraminidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-2 Fucosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-2Galβ1-4Glc-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Galactosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled</p>	Pass

Assay Name/Specification	Lot # 10084147
<p>α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (α1-3 Mannosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Xylosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Galactosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Mannosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 100 units of β1-3 Galactosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Mannosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 100 units of β1-3 Galactosidase</p>	Pass

Assay Name/Specification	Lot # 10084147
incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Alicia Bielik
Production Scientist
04 Nov 2020



Michael Tonello
Packaging Quality Control Inspector
04 Nov 2020