

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Bacteroides Heparinase I

Catalog Number: P0735L Concentration: 12,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will liberate 1.0

µmol unsaturated oligosaccharides from porcine mucosal heparin per

minute at 30°C and pH 7.0 in a total reaction volume of 100 μl.

Packaging Lot Number: 10109405 Expiration Date: 05/2022 -80°C Storage Temperature:

Storage Conditions: 100 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, 5 mM CaCl2, (pH 7.5 @ 25°C)

Specification Version: PS-P0735S/L v1.0

Bacteroides Heparinase I Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
P0735LVIAL	Bacteroides Heparinase I	10109404	Pass	
B0735SVIAL	Bacteroides Heparinase Reaction Buffer (10X)	10105362	Pass	

Assay Name/Specification	Lot # 10109405
Protein Purity Assay (SDS-PAGE) Bacteroides Heparinase I is ≥ 95% pure as determined by SDS-PAGE analysis using	Pass
Coomassie Blue detection.	_
Protease Activity (SDS-PAGE) A 20 μl reaction in 1X Heparinase Reaction Buffer containing 24 μg of a standard mixture of proteins and a minimum of 120 units of Bacteroides Heparinase I incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Sulfatase Activity (2-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled 2-O-Sulfatase substrate (ΔUA2S-(1-4)-GlcNS6S-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Sulfatase and Uronidase Activity (N,6-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled N,6-O-Sulfatase substrate (ΔUA-(1-4)-GlcNS6S-AMC) and 24 units	Pass



P0735L / Lot: 10109405

Page 1 of 3

Assay Name/Specification	Lot # 10109405
of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable	
activity as determined by thin layer chromatography.	
Changidana Antivity (84.3 Calastanidana)	Pass
Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of	Fass
fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC)	
and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in	
no detectable activity as determined by thin layer chromatography.	
	_
Glycosidase Activity (β1-4 Galactosidase)	Pass
A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC)	
and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in	
no detectable activity as determined by thin layer chromatography.	
Glycosidase Activity (β-N-Acetylgalactosaminidase)	Pass
A 10 µl reaction in Heparinase Reaction Buffer containing 1 nM of	
fluorescently-labeled β-N-Acetylgalactosaminidase substrate	
(GalNAcβ1-4Galβ1-4Glc-AMC) and 24 units of Bacteroides Heparinase I incubated for 20	
hours at 30°C results in no detectable activity as determined by thin layer chromatography.	
Chilomatography.	
Glycosidase Activity (β-N-Acetylglucosaminidase)	Pass
A 10 µl reaction in Heparinase Reaction Buffer containing 1 nM of	
fluorescently-labeled β-N-Acetylglucosaminidase substrate	
(GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 24 units of Bacteroides Heparinase I incubated	
for 20 hours at 30°C results in no detectable activity as determined by thin layer	
chromatography.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



P0735L / Lot: 10109405

Page 2 of 3

Alicia Bielik Production Scientist

17 Jun 2021

Michael Tonello
Packaging Quality Control Inspector

17 Jun 2021

