

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Bacteroides Heparinase III

Catalog Number: P0737S Concentration: 700 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will liberate 1.0

µmol unsaturated oligosaccharides from heparan sulfate per minute at

30°C and pH 7.0 in a total reaction volume of 100 μl.

Packaging Lot Number: 10070624
Expiration Date: 02/2021
Storage Temperature: -80°C

Storage Conditions: 100 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, 5 mM CaCl2, (pH 7.5 @ 25°C)

Specification Version: PS-P0737S/L v1.0

Bacteroides Heparinase III Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
P0737SVIAL	Bacteroides Heparinase III	10067395	Pass	
B0735SVIAL	Bacteroides Heparinase Reaction Buffer (10X)	10052584	Pass	

Assay Name/Specification	Lot # 10070624
Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 1 unit of Bacteroides Heparinase III incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 1 unit of Bacteroides Heparinase III incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 1 unit of Bacteroides Heparinase III incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass



P0737S / Lot: 10070624 Page 1 of 3

Assay Name/Specification	Lot # 10070624
Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 1 unit of Bacteroides Heparinase III incubated for 20 hours at 30° C results in no detectable activity as determined by thin layer chromatography.	Pass
Protease Activity (SDS-PAGE) A 20 µl reaction in 1X Heparinase Reaction Buffer containing 24 µg of a standard mixture of proteins and a minimum of 5 units of Bacteroides Heparinase III incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Protein Purity Assay (SDS-PAGE) Bacteroides Heparinase III is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Sulfatase Activity (2-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled 2-O-Sulfatase substrate (ΔUA2S-(1-4)-GlcNS6S-AMC) and 1 unit of Bacteroides Heparinase III incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Sulfatase and Uronidase Activity (N,6-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled N,6-O-Sulfatase substrate (ΔUA-(1-4)-GlcNS6S-AMC) and 1 unit of Bacteroides Heparinase III incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



P0737S / Lot: 10070624

Page 2 of 3

Alicia Bielik **Production Scientist**

30 Jul 2020

Josh Hersey Packaging Quality Control Inspector

30 Jul 2020

P0737S / Lot: 10070624

Page 3 of 3