

New England Biolabs Certificate of Analysis

Product Name: α 1-2, 3, 4, 6 Fucosidase
Catalog Number: P0748L
Concentration: 8,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to cleave > 95% of the α -L-fucose from 1 nmol of Fuc α 1-2Gal α 1-4Glc-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μ l.
Packaging Lot Number: 10055872
Expiration Date: 10/2020
Storage Temperature: 4°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0748S/L v2.0

α1-2, 3, 4, 6 Fucosidase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0748LVIAL	α 1-2, 3, 4, 6 Fucosidase	10055871	Pass
B9001SVIAL	Purified BSA	10014762	Pass
B1727SVIAL	10X GlycoBuffer 1	10041785	Pass

Assay Name/Specification	Lot # 10055872
Protease Activity (SDS-PAGE) A 20 μ l reaction in 1X Glyco Buffer 1 containing 24 μ g of a standard mixture of proteins and a minimum of 40 units of α 1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Protein Purity Assay (SDS-PAGE) α 1-2, 3, 4, 6 Fucosidase is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Glycosidase Activity (Endo F1, F2, H) A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 16 units of α 1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (Endo F2, F3)	Pass

Assay Name/Specification	Lot # 10055872
<p>A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (PNGase F) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Glucosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fuca1-2)Galβ1-4Glc-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Neuraminidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Galactosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Mannosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Galactosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled</p>	Pass

Assay Name/Specification	Lot # 10055872
<p>α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (α1-6 Mannosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Mannosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Xylosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 16 units of α1-2, 3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 16 units of α1-2,</p>	Pass

Assay Name/Specification	Lot # 10055872
3, 4, 6 Fucosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	

This product has been tested and shown to be in compliance with all specifications.



Tony Spear-Alfonso
Production Scientist
12 Sep 2018



Michael Tonello
Packaging Quality Control Inspector
04 Dec 2019