240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Endo F2
Catalog #: P0772S

Concentration: 8,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to cleave >95% of the carbohydrate from 10 µg Porcine Fibrinogen in 1

hour at 37°C in a total reaction volume of 10 µl.

 Lot #:
 0011612

 Assay Date:
 12/2016

 Expiration Date:
 12/2018

 Storage Temp:
 -20°C

Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)

Specification Version: PS-P0772S v1.0 Effective Date: 11 Nov 2016

Assay Name/Specification (minimum release criteria)	Lot #0011612
Glycosidase Activity (β-Mannosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β -Xylosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β -Xylosidase substrate (Xyl β 1-4Xyl β 1-4Xyl β 1-4Xyl-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β 1-3 Galactosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β -Galactosidase substrate (Gal β 1-3GlcNAc β 1-4Gal β 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β 1-4 Galactosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β -Galactosidase substrate (Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc -AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β -N-Acetylgalactosaminidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β -N-Acetylgalactosaminidase substrate (GalNAc β 1-4Gal β 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass









New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0011612
Glycosidase Activity (β -N-Acetylglucosaminidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled β -N-Acetylglucosaminidase substrate (GlcNAc β 1-4GlcNAc β 1-4GlcNAc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α -Glucosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Glucosidase substrate (Glc α 1-6Glc α 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-Neuraminidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α 1-2 Fucosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-2Gal β 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α 1-3 Fucosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity ($\alpha 1$ -3 Galactosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal $\alpha 1$ -3Gal $\beta 1$ -4GlcNAc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α 1-3 Mannosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-3Man β 1-4GlcNAc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α 1-6 Galactosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-6Gal α 1-6Glc α 1-2Fru-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α 1-6 Mannosidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-6(Man α 1-3)Man-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α -N-Acetylgalactosaminidase) - A 10 μ l reaction in Glyco Buffer 4 containing 1 nM of fluorescently-labeled α -N-Acetylgalactosaminidase substrate (GalNAc α 1-3(Fuc α 1-2)Gal β 1-4Glc-AMC) and 16 units of Endo F2 incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0011612
Protease Activity (SDS-PAGE) - A 20 μl reaction in 1X Glyco Buffer 4 containing 24 μg of a standard mixture of proteins and a minimum of 40 units of Endo F2 incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue.	Pass
Protein Purity Assay (SDS-PAGE) - Endo F2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

Authorized by Derek Robinson 11 Nov 2016







Inspected by Alicia Bielik 20 Dec 2016