

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Proteinase K, Molecular Biology Grade

Catalog Number: P8107S Concentration: 800 U/ml

Unit Definition: One unit will digest urea-denatured hemoglobin at 37°C (pH 7.5) per

minute to produce equal absorbance as 1.0 μmol L-tyrosine using

Folin & Ciocalteu's phenol reagent.

Packaging Lot Number: 10130705 Expiration Date: 11/2024 Storage Temperature: -20°C

Storage Conditions: 20 mM Tris-HCl, 1 mM CaCl2, 50% Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-P8107S v2.0

Proteinase K, Molecular Biology Grade Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
P8107SVIAL	Proteinase K, Molecular Biology Grade	10121120	Pass	

Assay Name/Specification	Lot # 10130705
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 8 units of Proteinase K, Molecular Biology Grade incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 RF I DNA and a minimum of 0.8 units of Proteinase K, Molecular Biology Grade incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 0.8 units of Proteinase K, Molecular Biology Grade incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
qPCR DNA Contamination (Eukaryotic Genomic) A minimum of 1.6 units of Proteinase K, Molecular Biology Grade is screened for the	Pass



P8107S / Lot: 10130705

Page 1 of 2

Assay Name/Specification	Lot # 10130705
presence of eukaryotic genomic DNA using SYBR® Green qPCR with universal primers for the 18S rRNA locus. Results are quantified using a standard curve generated from purified E. album genomic DNA. The measured level of eukaryotic genomic DNA contamination is ≤ 2.5 pg DNA/µI.	
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of fluorescein labeled RNA transcript and a minimum of 0.8 units of Proteinase K, Molecular Biology Grade is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 4 units of Proteinase K, Molecular Biology Grade incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Alicia Bielik Production Scientist 23 Nov 2021 Michael Tonello

Packaging Quality Control Inspector

23 Nov 2021

P8107S / Lot: 10130705